

Comment on “Exact results for survival probability in the multistate Landau-Zener model”

B. E. Dobrescu

*Department of Physics, Texas A&M University,
College Station, TX 77843-4242, USA*

N. A. Sinitsyn

Department of Physics, University of Texas at Austin, Austin TX 78712-1081, USA

Abstract

We correct the proof of Brundobler-Elser formula (BEF) provided in [2004 *J. Phys. B: At. Mol. Opt. Phys.* **37** 4069] and continued in Appendix of [2005 *J. Phys. B: At. Mol. Opt. Phys.* **38** 907]. After showing that some changes of variables employed in these articles are used erroneously, we propose an alternative change of variables which solves the problem. In our proof, we reveal the connection between the BEF for a general N -level Landau-Zener system and the exactly solvable bow-tie model. The special importance of the diabatic levels with maximum/minimum slope is emphasized throughout.

I. INTRODUCTION

In Ref. [1] Volkov and Ostrovsky (referred to below as V-O) proposed a rigorous mathematical proof of the Brundobler-Elser formula (BEF) [3] for a general N -level Landau-Zener (LZ) system. The approach of [1] was to consider a formal solution in terms of an infinite series in powers of the coupling constants, with further calculation of every term and summation of the resulting algebraic series. Following this strategy, these authors have demonstrated the exact cancellations of some quantum paths and have reduced the problem to the calculation of a sub-series (5.1) (the numbering of formulas is as in [1]).

In the calculation of this remaining sub-series (5.1) in [1] V-O made the error of assuming a wrong domain of integration for their variables x_j (see Eq. (5.2) of [1]).

In Ref. [2], whose main text is dedicated to a different subject, V-O included an Appendix in which a new attempt is made at calculating the sub-series (5.1). After acknowledging the error made in [1], V-O proceeded to employ a different change of variables, previously proposed by Kayanuma and Fukuchi in [5].

We would like to point out that this derivation contained in the Appendix of [2] is still erroneous. After Eq. (A.4) (in the notation of Ref. [2]) the authors claim that the integrand is symmetrical under the permutation of variables $x_0, \dots, x_{m-1}$. It is, however, straightforward to check, by direct examination of the terms containing the 4-th power in the coupling constants V_i , that this is not true; nor is it symmetrical under exchange of pairs of variables $(x_0, y_0), \dots, (x_{m-1}, y_{m-1})$.

The following example will support these considerations: the term corresponding to $l = m = 2$ in Eq. (5.1) of [1] reads

$$\begin{aligned} & \sum_{k_1 \neq 1}^N |V_{1k_1}|^2 \sum_{k_2 \neq 1}^N |V_{1k_2}|^2 \int_{-\infty}^{\infty} dt_4 \int_{-\infty}^{t_4} dt_3 \int_{-\infty}^{t_3} dt_2 \int_{-\infty}^{t_2} dt_1 \\ & \times \exp [i (\varepsilon_1 - \varepsilon_{k_1}) (t_4 - t_3) + i (\varepsilon_1 - \varepsilon_{k_2}) (t_2 - t_1)] \\ & \times \exp \left[\frac{i}{2} (\beta_1 - \beta_{k_1}) (t_4^2 - t_3^2) + \frac{i}{2} (\beta_1 - \beta_{k_2}) (t_2^2 - t_1^2) \right]. \end{aligned} \quad (1)$$

Let us consider the simplest model beyond the two-level LZ system, namely the one with $N = 3$. Then, the sum above has 4 terms

$$\begin{aligned} I_{22} = & |V_{12}|^4 \int_{-\infty}^{\infty} dt_4 \int_{-\infty}^{t_4} dt_3 \int_{-\infty}^{t_3} dt_2 \int_{-\infty}^{t_2} dt_1 \\ & \times \exp \left[i (\varepsilon_1 - \varepsilon_2) (t_4 - t_3 + t_2 - t_1) + \frac{i}{2} (\beta_1 - \beta_2) (t_4^2 - t_3^2 + t_2^2 - t_1^2) \right], \end{aligned} \quad (2)$$

$$I_{33} = |V_{13}|^4 \int_{-\infty}^{\infty} dt_4 \int_{-\infty}^{t_4} dt_3 \int_{-\infty}^{t_3} dt_2 \int_{-\infty}^{t_2} dt_1 \quad (3)$$

$$\times \exp \left[i (\varepsilon_1 - \varepsilon_3) (t_4 - t_3 + t_2 - t_1) + \frac{i}{2} (\beta_1 - \beta_3) (t_4^2 - t_3^2 + t_2^2 - t_1^2) \right],$$

$$I_{23} = |V_{12}|^2 |V_{13}|^2 \int_{-\infty}^{\infty} dt_4 \int_{-\infty}^{t_4} dt_3 \int_{-\infty}^{t_3} dt_2 \int_{-\infty}^{t_2} dt_1 \quad (4)$$

$$\times \exp [i (\varepsilon_1 - \varepsilon_2) (t_4 - t_3) + i (\varepsilon_1 - \varepsilon_3) (t_2 - t_1)]$$

$$\times \exp \left[\frac{i}{2} (\beta_1 - \beta_2) (t_4^2 - t_3^2) + \frac{i}{2} (\beta_1 - \beta_3) (t_2^2 - t_1^2) \right],$$

$$I_{32} = |V_{13}|^2 |V_{12}|^2 \int_{-\infty}^{\infty} dt_4 \int_{-\infty}^{t_4} dt_3 \int_{-\infty}^{t_3} dt_2 \int_{-\infty}^{t_2} dt_1 \quad (5)$$

$$\times \exp [i (\varepsilon_1 - \varepsilon_3) (t_4 - t_3) + i (\varepsilon_1 - \varepsilon_2) (t_2 - t_1)]$$

$$\times \exp \left[\frac{i}{2} (\beta_1 - \beta_3) (t_4^2 - t_3^2) + \frac{i}{2} (\beta_1 - \beta_2) (t_2^2 - t_1^2) \right].$$

The change of variables [5] used by V-O in Appendix of [2] amounts to

$$t_1 = x_1, \quad x_1 \in (-\infty, \infty)$$

$$t_2 = x_1 + y_1, \quad y_1 \in [0, \infty)$$

$$t_3 = x_2 + y_1, \quad x_2 \in [x_1, \infty)$$

$$t_4 = x_2 + y_1 + y_2, \quad y_2 \in [0, \infty)$$

The integrals I_{22} and I_{33} can be easily evaluated (see [5]), e.g.

$$I_{22} \equiv |V_{12}|^4 \int_{-\infty}^{\infty} dx_2 \int_{-\infty}^{x_2} dx_1 \int_0^{\infty} dy_2 \int_0^{\infty} dy_1 \quad (6)$$

$$\times \exp \left[i (\varepsilon_1 - \varepsilon_2) (y_1 + y_2) + \frac{i}{2} (\beta_1 - \beta_2) (y_1 + y_2)^2 + i (\beta_1 - \beta_2) (y_1 x_1 + y_2 x_2) \right]$$

$$= |V_{12}|^4 \frac{1}{2} \int_{-\infty}^{\infty} dx_2 \int_{-\infty}^{\infty} dx_1 \int_0^{\infty} dy_2 \int_0^{\infty} dy_1 \times$$

$$\times \exp \left[i (\varepsilon_1 - \varepsilon_2) (y_1 + y_2) + \frac{i}{2} (\beta_1 - \beta_2) (y_1 + y_2)^2 + i (\beta_1 - \beta_2) (y_1 x_1 + y_2 x_2) \right]$$

$$= |V_{12}|^4 \frac{1}{2} \int_0^{\infty} dy_2 \int_0^{\infty} dy_1 \frac{2\pi}{|\beta_1 - \beta_2|} \delta(y_1) \frac{2\pi}{|\beta_1 - \beta_2|} \delta(y_2) \times$$

$$\times \exp \left[i (\varepsilon_1 - \varepsilon_2) (y_1 + y_2) + \frac{i}{2} (\beta_1 - \beta_2) (y_1 + y_2)^2 \right] = \frac{|V_{12}|^4}{2} \left(\frac{\pi}{\beta_1 - \beta_2} \right)^2$$

and, similarly, $I_{33} = \frac{|V_{13}|^4}{2} \left(\frac{\pi}{\beta_1 - \beta_3} \right)^2$.

The sum of the remaining two integrals reads

$$\begin{aligned}
I_{23} + I_{32} = & |V_{12}|^2 |V_{13}|^2 \int_{-\infty}^{\infty} dx_1 \int_{x_1}^{\infty} dx_2 \int_0^{\infty} dy_2 \int_0^{\infty} dy_1 \\
& \times \left\{ \exp [i (\varepsilon_1 - \varepsilon_2) y_2 + i (\varepsilon_1 - \varepsilon_3) y_1] \right. \\
& \times \exp \left[\frac{i}{2} (\beta_1 - \beta_2) (2x_2 y_2 + y_2^2) + \frac{i}{2} (\beta_1 - \beta_3) (2x_1 y_1 + y_1^2) + i (\beta_1 - \beta_2) y_1 y_2 \right] \\
& + \exp [i (\varepsilon_1 - \varepsilon_3) y_2 + i (\varepsilon_1 - \varepsilon_2) y_1] \\
& \left. \times \exp \left[\frac{i}{2} (\beta_1 - \beta_3) (2x_2 y_2 + y_2^2) + \frac{i}{2} (\beta_1 - \beta_2) (2x_1 y_1 + y_1^2) + i (\beta_1 - \beta_3) y_1 y_2 \right] \right\}.
\end{aligned} \tag{7}$$

Contrary to the claims made by V-O after Eq. (A.4) of [2], the integrand of (7) is clearly *not symmetric* with respect to interchanging the pairs (x_1, y_1) and (x_2, y_2) . Indeed, the terms $(\beta_1 - \beta_2) y_1 y_2 \neq (\beta_1 - \beta_3) y_1 y_2$ belong to different terms of the sum in the integrand and are left unchanged by the permutation. Therefore, one cannot extend the domain of integration of variable x_2 to $(-\infty, +\infty)$ as proposed by V-O in Eq. (A.5) of [2].

While the mathematical errors made by V-O are presented above in detail only for completeness, there is also a more general argument as to why the calculation of the series (5.1) in both Ref. [1] and Ref. [2] is erroneous. One can notice that this series represents a formal solution for a special case of the multi-state LZ-model, in which all states interact with only one special level with slope β_1 . Series (5.1) is the probability amplitude for the system to remain on this special level after all crossings if only this level has been initially populated. This observation does *not* assume at all that the slope β_1 is an extremum. Therefore, if the proof provided by V-O in [1] and [2] were correct, then their summation of the series (5.1) that does *not* use the assumption of a maximal/minimal slope for the level 1 could be applied to such a LZ-model with an *arbitrary* slope β_1 .

However, it is well known from numerical simulations [3] of this model that if β_1 is *not* an extremum, then the BEF does *not* hold, and the transition probability for $1 \rightarrow 1$ depends essentially on the parameters ε_i . Also, the exactly solvable case of the above example in which all $\varepsilon_i = 0$, the bow-tie model [4], proves that the diagonal elements of the S matrix are *different* from BEF *except* for the levels with *maximal/minimal* slope.

The total ignorance of this very important property of the slope β_1 being an *extremum* in the calculation of series (5.1) by V-O has actually been the trigger that prompted our search for their mathematical mistakes.

II. PROOF OF BEF

We will split the proof of BEF in three steps. The first step has been correctly performed in [1] and we only briefly mention here the main results in order to introduce the notation and as a starting point for the next steps.

It has been shown in [1] that *if* the slope β_1 is *maximal/minimal* then the transition amplitude for the transition $1 \rightarrow 1$ has the form:

$$\begin{aligned}
S_{11} \equiv \langle 1 | \hat{U}(\infty, -\infty) | 1 \rangle &= 1 + \sum_{m=1}^{\infty} (-1)^m \sum_{k_1=2}^N \sum_{k_2=2}^N \cdots \sum_{k_m=2}^N |V_{1k_1}|^2 |V_{1k_2}|^2 \cdots |V_{1k_m}|^2 \times \\
&\times \int_{-\infty}^{\infty} d\tau_1 \int_{-\infty}^{\tau_1} d\tau_2 \cdots \int_{-\infty}^{\tau_{2m-1}} d\tau_{2m} \\
&\times \exp \left[i \frac{B(k_1)}{2} (\tau_1^2 - \tau_2^2) + i \frac{B(k_2)}{2} (\tau_3^2 - \tau_4^2) + \cdots + i \frac{B(k_m)}{2} (\tau_{2m-1}^2 - \tau_{2m}^2) \right] \\
&\times \exp [iE(k_1)(\tau_1 - \tau_2) + iE(k_2)(\tau_3 - \tau_4) + \cdots + iE(k_m)(\tau_{2m-1} - \tau_{2m})], \quad (8)
\end{aligned}$$

where

$$B(k_j) \equiv \beta_1 - \beta_{k_j} \quad \text{and} \quad E(k_j) \equiv \varepsilon_1 - \varepsilon_{k_j}, \quad j = 1, 2, \dots, m. \quad (9)$$

$\hat{U}$ is the time-evolution operator and N is the number of levels in the system.

Series (8) is equivalent to the series (5.1) in [1]. From this point on, Volkov and Ostrovsky do *not* use the property of β_1 being *maximal/minimal* anymore in [1] and in the Appendix of [2], and the changes of variables they employ are used erroneously, as shown in the Introduction.

In the second step we prove that *if* β_1 is *maximal/minimal*, then $S_{11} \equiv \langle 1 | \hat{U}(\infty, -\infty) | 1 \rangle$ does *not* depend on $E(l_j)$, for any $l_j = 2, 3, \dots, N$ and any $j = 1, 2, \dots, m$.

From Eq.(8) it follows that:

$$\begin{aligned}
\frac{\partial S_{11}}{\partial E(l_j)} &= \sum_{m=1}^{\infty} (-1)^m \sum_{k_1=2}^N \sum_{k_2=2}^N \cdots \sum_{k_m=2}^N |V_{1k_1}|^2 |V_{1k_2}|^2 \cdots |V_{1k_m}|^2 \times \\
&\times \int_{-\infty}^{\infty} d\tau_1 \int_{-\infty}^{\tau_1} d\tau_2 \cdots \int_{-\infty}^{\tau_{2m-1}} d\tau_{2m} \\
&\times i [\delta(l_j, k_1)(\tau_1 - \tau_2) + \delta(l_j, k_2)(\tau_3 - \tau_4) + \cdots + \delta(l_j, k_m)(\tau_{2m-1} - \tau_{2m})] \\
&\times \exp \left[i \frac{B(k_1)}{2} (\tau_1^2 - \tau_2^2) + i \frac{B(k_2)}{2} (\tau_3^2 - \tau_4^2) + \cdots + i \frac{B(k_m)}{2} (\tau_{2m-1}^2 - \tau_{2m}^2) \right] \\
&\times \exp [iE(k_1)(\tau_1 - \tau_2) + iE(k_2)(\tau_3 - \tau_4) + \cdots + iE(k_m)(\tau_{2m-1} - \tau_{2m})], \quad (10)
\end{aligned}$$

where $\delta(l_j, k_p)$ is the Kronecker delta.

Next, we introduce the well-known change of variables (see, e.g., textbook [7], page 114):

$$\begin{aligned}\tau_1 &= x_1 \in (-\infty, \infty) \\ \tau_{j+1} &= \tau_j - x_{j+1}, \quad \text{with } x_{j+1} \in [0, \infty), \quad j = 1, 2, \dots, 2m-1.\end{aligned}\tag{11}$$

From Eqs. (10) and (11) it follows that

$$\begin{aligned}\frac{\partial S_{11}}{\partial E(l_j)} &= \sum_{m=1}^{\infty} (-1)^m \sum_{k_1=2}^N \sum_{k_2=2}^N \cdots \sum_{k_m=2}^N |V_{1k_1}|^2 |V_{1k_2}|^2 \cdots |V_{1k_m}|^2 \times \\ &\times \int_{-\infty}^{\infty} dx_1 \int_0^{\infty} dx_2 \int_0^{\infty} dx_3 \cdots \int_0^{\infty} dx_{2m} \\ &\times i [\delta(l_j, k_1) x_2 + \delta(l_j, k_2) x_4 + \cdots + \delta(l_j, k_m) x_{2m}] \\ &\times F(x_1, x_2, x_3, \dots, x_{2m}),\end{aligned}\tag{12}$$

where

$$\begin{aligned}F(x_1, x_2, x_3, \dots, x_{2m}) &= \exp \left[-\frac{i}{2} G(x_2, x_4, \dots, x_{2m-2}, x_{2m}) \right] \times \\ &\times \exp [i \{B(k_1) x_2 + B(k_2) x_4 + \cdots + B(k_m) x_{2m}\} x_1] \\ &\times \exp [-i \{B(k_2) x_4 + B(k_3) x_6 + \cdots + B(k_m) x_{2m}\} x_3] \\ &\times \exp [-i \{B(k_3) x_6 + B(k_4) x_8 + \cdots + B(k_m) x_{2m}\} x_5] \\ &\vdots \\ &\times \exp [-i \{B(k_m) x_{2m}\} x_{2m-1}],\end{aligned}\tag{13}$$

and

$$\begin{aligned}G(x_2, x_4, \dots, x_{2m-2}, x_{2m}) &= B(k_1) (x_2^2) + B(k_2) (2x_4 x_2 + x_4^2) \\ &+ B(k_3) (2x_6 x_2 + 2x_6 x_4 + x_6^2) \\ &\vdots \\ &+ B(k_m) (2x_{2m} x_2 + 2x_{2m} x_4 + \cdots + 2x_{2m} x_{2m-2} + x_{2m}^2) \\ &- 2 [E(k_1) x_2 + E(k_2) x_4 + \cdots + E(k_m) x_{2m}].\end{aligned}\tag{14}$$

Upon integrating over x_1 in Eq.(12) one obtains:

$$\begin{aligned}\frac{\partial S_{11}}{\partial E(l_j)} &= \sum_{m=1}^{\infty} (-1)^m \sum_{k_1=2}^N \sum_{k_2=2}^N \cdots \sum_{k_m=2}^N |V_{1k_1}|^2 |V_{1k_2}|^2 \cdots |V_{1k_m}|^2 \times \\ &\times \int_0^{\infty} dx_2 \int_0^{\infty} dx_3 \cdots \int_0^{\infty} dx_{2m}\end{aligned}$$

$$\begin{aligned}
& \times i [\delta(l_j, k_1) x_2 + \delta(l_j, k_2) x_4 + \cdots + \delta(l_j, k_m) x_{2m}] \\
& \times \exp \left[-\frac{i}{2} G(x_2, x_4, \dots, x_{2m-2}, x_{2m}) \right] \\
& \times 2\pi \delta(B(k_1) x_2 + B(k_2) x_4 + \cdots + B(k_m) x_{2m}) \\
& \times \exp[-i \{B(k_2) x_4 + B(k_3) x_6 + \cdots + B(k_m) x_{2m}\} x_3] \\
& \times \exp[-i \{B(k_3) x_6 + B(k_4) x_8 + \cdots + B(k_m) x_{2m}\} x_5] \\
& \vdots \\
& \times \exp[-i \{B(k_m) x_{2m}\} x_{2m-1}].
\end{aligned} \tag{15}$$

From Eq.(9) it follows that, for any k_j , $B(k_j) > 0$ if β_1 is maximal, and $B(k_j) < 0$ if β_1 is minimal. Therefore, $\delta(B(k_1) x_2 + B(k_2) x_4 + \cdots + B(k_m) x_{2m}) = 0$ unless $x_2 = x_4 = \cdots = x_{2m} = 0$. The presence of the term

$$[\delta(l_j, k_1) x_2 + \delta(l_j, k_2) x_4 + \cdots + \delta(l_j, k_m) x_{2m}] \tag{16}$$

in the integrands of Eq.(15) ensures that each of the integrals is zero.

This argument can be made rigorous by regularizing the behavior of integrals at $\pm\infty$, as follows

$$\begin{aligned}
I_q & \equiv \int_{-\infty}^{\infty} dx_1 \int_0^{\infty} dx_2 \int_0^{\infty} dx_3 \cdots \int_0^{\infty} dx_{2m} F(x_1, \dots, x_{2m}) x_{2q} \\
& = \lim_{\eta \rightarrow 0_+} \left\{ \int_0^{\infty} dx_2 \int_0^{\infty} dx_4 \cdots \int_0^{\infty} dx_{2m} \int_0^{\infty} dx_1 \int_0^{\infty} dx_3 \cdots \int_0^{\infty} dx_{2m-1} \times \right. \\
& \quad \times x_{2q} F(x_1, \dots, x_{2m}) e^{-\eta(x_1+x_3+\cdots+x_{2m-1})} \\
& \quad + \int_0^{\infty} dx_2 \int_0^{\infty} dx_4 \cdots \int_0^{\infty} dx_{2m} \int_{-\infty}^0 dx_1 \int_0^{\infty} dx_3 \cdots \int_0^{\infty} dx_{2m-1} \times \\
& \quad \times x_{2q} F(x_1, \dots, x_{2m}) e^{-\eta(-x_1+x_3+\cdots+x_{2m-1})} \left. \right\},
\end{aligned} \tag{17}$$

for any $q = 1, 2, \dots, m$. Upon integrating over the x_j with odd j in Eq.(17), one obtains

$$I_q = \lim_{\eta \rightarrow 0_+} \int_0^{\infty} dx_2 \int_0^{\infty} dx_4 \cdots \int_0^{\infty} dx_{2m} W(x_2, x_4, \dots, x_{2m}; \eta), \tag{18}$$

where

$$\begin{aligned}
W(x_2, x_4, \dots, x_{2m}; \eta) & = \exp \left[-\frac{i}{2} G(x_2, x_4, \dots, x_{2m}) \right] \times \\
& \quad \times \frac{2\eta x_{2q}}{[B(k_1) x_2 + B(k_2) x_4 + \cdots + B(k_m) x_{2m}]^2 + \eta^2} \\
& \quad \times \frac{1}{i [B(k_2) x_4 + B(k_3) x_6 + \cdots + B(k_m) x_{2m}] + \eta}
\end{aligned}$$

$$\begin{aligned} & \vdots \\ & \times \frac{1}{i [B(k_m) x_{2m}] + \eta}. \end{aligned} \quad (19)$$

Since either $B(k_j) > 0$ or $B(k_j) < 0$ for any k_j , from Eqs. (18) and (19) it follows that

$$\begin{aligned} |I_q| & \leq \lim_{\eta \rightarrow 0_+} \int_0^\infty dx_2 \int_0^\infty dx_4 \cdots \int_0^\infty dx_{2m} |W(x_2, x_4, \dots, x_{2m}; \eta)| \\ & = \lim_{\eta \rightarrow 0_+} \int_0^{\eta/|B(k_1)|} dx_2 \int_0^{\eta/|B(k_2)|} dx_4 \cdots \int_0^{\eta/|B(k_m)|} dx_{2m} \times \\ & \quad \times \frac{2\eta x_{2q}}{[B(k_1)x_2 + B(k_2)x_4 + \cdots + B(k_m)x_{2m}]^2 + \eta^2} \\ & \quad \times \frac{1}{\sqrt{[B(k_2)x_4 + B(k_3)x_6 + \cdots + B(k_m)x_{2m}]^2 + \eta^2}} \cdots \frac{1}{\sqrt{[B(k_m)x_{2m}]^2 + \eta^2}}. \end{aligned} \quad (20)$$

For any continuous function f defined on a compact interval $[a, b]$ the relation $\int_a^b f(x) dx = f(\xi)(b-a)$ holds, for some $\xi \in [a, b]$. Hence, Eq.(20) reduces to

$$\begin{aligned} |I_q| & \leq \frac{1}{|B(k_1)B(k_2) \cdots B(k_m)|} \lim_{\eta \rightarrow 0_+} 2\eta \frac{\xi_q}{|B(k_q)|} \frac{1}{(\xi_1 + \xi_2 + \cdots + \xi_m)^2 + 1} \times \\ & \quad \times \frac{1}{\sqrt{(\xi_2 + \xi_3 + \cdots + \xi_m)^2 + 1}} \cdots \frac{1}{\sqrt{\xi_m^2 + 1}} = 0, \end{aligned} \quad (21)$$

where $\xi_j \in [0, 1]$, for any $j = 1, 2, \dots, m$. Eq.(21) shows that $I_q = 0$ for any $q = 1, 2, \dots, m$, and consequently $\frac{\partial S_{11}}{\partial E(l_j)} = 0$ for any $E(l_j)$.

The fact that β_1 is *maximal/minimal* has played a key role in the arguments above. These arguments fail to hold if β_1 is not an extremum, since then the argument of $\delta(B(k_1)x_2 + B(k_2)x_4 + \cdots + B(k_m)x_{2m})$ would have other zeros, besides the obvious $x_2 = x_4 = \cdots = x_{2m} = 0$.

In the last step, we reveal the connection between the BEF for a general N -level LZ-system and the exactly solvable bow-tie model [4], in which all levels interact with only one special level (SL). Indeed, since S_{11} does not depend on $E(l_j)$ if β_1 is *maximal/minimal*, we can safely set $E(l_j) = 0$ for any $l_j = 2, 3, \dots, N$. The form for S_{11} (Eq.(8)) with all $E(l_j) = 0$ is exactly what one obtains for a bow-tie model if the SL has slope β_1 , and a perturbation expansion is being developed for it.

Since the Brundobler-Elser conjecture for the bow-tie model was proven analytically by Ostrovsky and Nakamura in [4], we only cite their statement from page 6947 of [4]: “*This hypothesis is confirmed within the present model.*”, and refer to their work for further details.

This ends the proof of the Brundobler-Elser formula for a general N -level LZ-system. Before drawing the conclusions, it is worthwhile to mention that, preceding [1], an alternative proof of BEF, based on analytic continuation in imaginary time, was proposed by Shytov in [6].

III. CONCLUSIONS

In summary, we revealed the deficiencies of the BEF proof proposed by Volkov and Ostrovsky in [1] and Appendix of [2]. We emphasized that using the important property of the slope β_1 being an extremum *only* to arrive at the sub-series (5.1) in their proof is *not enough*, regardless of the changes of variables they attempted, and this property still has to play a crucial role in evaluating (5.1).

We have constructed a proof of BEF that corrects for this shortcoming. Indeed, in our proof, the fact that the slope β_1 is maximal/minimal had to be used for three different purposes: 1) to arrive at Eq.(5.1) as was shown in [1] ; 2) to prove that S_{11} does not depend on $E(l_j)$; 3) to employ the fact that the Brundobler-Elser conjecture for maximum/minimum slope was proved for the particular case of the bow-tie model [4].

Note added.-After this work was submitted to publication we received a Reply by Volkov and Ostrovsky [8] in which these authors now verify, by direct calculation, the important point on which they missed before in [1] and Appendix of [2], and which we have advocated throughout our Comment, namely that in order for the BEF to hold the important property of β_1 being an *extreme slope (ES)* must be used manifestly in the evaluation of *series* (5.1).

In their Reply V-O abandon the old changes of variables and use this time the same well-known change of variables (11) (see, e.g., textbook [7], page 114) as we have employed. We note that this change of variables, while previously used by V-O in a different context (e.g., in the proof of the no-go theorem [2]), was never employed in their attempts of calculating *series* (5.1) in [1] and Appendix of [2]. The point at issue in our Comment is, however, not about what changes of variables should be or not be used in the proof of BEF. Mathematical inconsistencies aside, it was definitely not the V-O's choice of integration variables (in [1] and Appendix of [2]) that prompted our Comment, but their *total neglect* of the very important *ES*-property of β_1 in evaluating *series* (5.1).

Though only the evaluation of 4-th order terms is presented in [8], our claim that one can set $\varepsilon_j = 0$ *only if* β_1 is an *ES* is reconfirmed, together with the fact that this *ES*-property must *still* be used even after setting $\varepsilon_j = 0$ in order to prove the Brundobler-Elser conjecture.

IV. ACKNOWLEDGMENTS

We thank V. L. Pokrovsky for encouragement. This work was supported by DOE grants DE-FG03-96ER45598 and DE-FG03-02ER45958, by NSF under the grants DMR-0115947 and DMR-0321572, and by the Welch Foundation.

-
- [1] Volkov M V and Ostrovsky V N 2004 *J. Phys. B: At. Mol. Opt. Phys.* **37** 4069-84
 - [2] Volkov M V and Ostrovsky V N 2005 *J. Phys. B: At. Mol. Opt. Phys.* **38** 907-15
 - [3] Brundobler S and Elser V 1993 *J. Phys. A: Math. Gen.* **26** 1211-27
 - [4] Ostrovsky V N and Nakamura H 1997 *J. Phys. A: Math. Gen.* **30** 6939-50
 - [5] Kayanuma Y and Fukuchi S 1985 *J. Phys. B: At. Mol. Phys.* **18** 4089-4093
 - [6] Shytov A V 2004 *Phys. Rev. A* **70** 052708
 - [7] Fetter A L and Walecka J D 1971 *Quantum Theory of Many-Particle Systems* (McGraw-Hill, New York)
 - [8] Volkov M V and Ostrovsky V N 2005 *J. Phys. B: At. Mol. Opt. Phys.*, this volume